

Bounds on 4D Conformal and Superconformal Field Theories

David Poland

Harvard University

March 21, 2011

(with David Simmons-Duffin [arXiv:1009.2087])

Motivation

- ▶ Conformal dynamics in 4D could play a role in BSM physics!
 - ▶ Walking/Conformal Technicolor [Holdom '81; ...]
 - ▶ Warped Extra Dimensions [Randall, Sundrum '99; ...]
 - ▶ Conformal Sequestering [Luty, Sundrum '01; Schmaltz, Sundrum '06]
 - ▶ Solution to $\mu/B\mu$ problem [Roy, Schmaltz '07; Murayama, Nomura, DP '07]
 - ▶ Flavor Hierarchies [Georgi, Nelson, Manohar '83; Nelson, Strassler '00]
- ▶ Ideas often make strong assumptions about operator dim's
 - ▶ E.g., Conf. Technicolor: Want $\dim H^\dagger H \gtrsim 4$ but $\dim H \sim 1$
- ▶ But it's hard to calculate *anything* in non-SUSY 4D CFTs!
- ▶ In $\mathcal{N} = 1$ SCFTs, we actually know lots about chiral operators, but not much about non-chiral operators...

Example: Nelson-Strassler Flavor Models ['00]

- ▶ Idea: Matter fields T_i have large anomalous dimensions γ_i under some CFT, flavor hierarchies generated dynamically!

$$W = T_1 \mathcal{O}_1 + T_2 \mathcal{O}_2 + y^{ij} T_i T_j H + \dots$$

- ▶ Interactions of matter T_i with CFT operators \mathcal{O}_i are marginal
- ▶ Yukawa couplings y^{ij} flow to zero at rate controlled by γ_i

$$y^{ij} T_i T_j H \rightarrow \left(\frac{\mu}{\Lambda} \right)^{\gamma_i + \gamma_j} y^{ij} T_i T_j H$$

- ▶ Since T_i are chiral, $\dim T_i = \frac{3}{2} R_{T_i}$ (superconformal $U(1)_R$)
- ▶ Can write down lots of concrete models and then *calculate* dimensions using a-maximization! [DP, Simmons-Duffin '09]

Example: Nelson-Strassler Flavor Models ['00]

- ▶ Soft-mass operators $K \sim \frac{1}{M_{pl}^2} X^\dagger X T_i^\dagger T_j$ also flow to zero
 - ▶ Rate controlled by $\dim T_i^\dagger T_j$
- ▶ Maybe can solve SUSY flavor problem???
- ▶ No way to calculate dimensions...
- ▶ Similar issue arises in Conformal Sequestering, $\mu/B\mu$ solution
- ▶ Can we say *anything* about $\dim T^\dagger T$, given $\dim T$?

Example: Nelson-Strassler Flavor Models ['00]

- ▶ Soft-mass operators $K \sim \frac{1}{M_{pl}^2} X^\dagger X T_i^\dagger T_j$ also flow to zero
 - ▶ Rate controlled by $\dim T_i^\dagger T_j$
- ▶ Maybe can solve SUSY flavor problem???
 - ▶ No way to calculate dimensions...
- ▶ Similar issue arises in Conformal Sequestering, $\mu/B\mu$ solution
- ▶ Can we say *anything* about $\dim T^\dagger T$, given $\dim T$?

- ▶ Recently, the papers:
 - [Rattazzi, Rychkov, Tonni, Vichi \[arXiv:0807.0004\]](#)
 - [Rychkov, Vichi \[arXiv:0905.2211\]](#)

addressed a similar question in non-SUSY CFTs, deriving *bounds* on $\dim \phi^2$ as a function of $\dim \phi$...

Outline

- 1 CFT Review
- 2 Bounds from Crossing Relations
- 3 Superconformal Blocks
- 4 Bounds on CFTs and SCFTs
- 5 Outlook

CFT Review: Primary Operators

- ▶ In addition to Poincaré generators P^a and M^{ab} , CFTs have dilatations D and special conformal generators K^a

$$[K^a, P^b] = 2\eta^{ab}D - 2M^{ab}$$

- ▶ *Primary* operators $\mathcal{O}^I(0)$ are defined by

$$[K^a, \mathcal{O}^I(0)] = 0$$

(descendants obtained by acting with P^a)

CFT Review: Primary Operators

- ▶ Primary 2-pt and 3-pt functions fixed by conformal symmetry in terms of dimensions and spins, up to overall coefficients $\lambda_{\mathcal{O}}$

$$\langle \mathcal{O}^{a_1 \dots a_l}(x_1) \mathcal{O}^{b_1 \dots b_l}(x_2) \rangle = \frac{I^{a_1 b_1} \dots I^{a_l b_l}}{x_{12}^{2\Delta}}$$

$$\langle \phi(x_1) \phi(x_2) \mathcal{O}^{a_1 \dots a_l}(x_3) \rangle = \frac{\lambda_{\mathcal{O}}}{x_{12}^{2d-\Delta+l} x_{23}^{\Delta-l} x_{13}^{\Delta-l}} Z^{a_1} \dots Z^{a_l}$$

$$\left(I^{ab} = \eta^{ab} - 2 \frac{x_{12}^a x_{12}^b}{x_{12}^2}, \quad Z^a = \frac{x_{31}^a}{x_{31}^2} - \frac{x_{32}^a}{x_{32}^2} \right)$$

- ▶ Higher n -pt functions *not* fixed by conformal symmetry alone, but are determined once spectrum and $\lambda_{\mathcal{O}}$'s are known...

CFT Review: Operator Product Expansion

Let ϕ be a scalar primary of dimension d in a 4D CFT:

$$\phi(x)\phi(0) = \sum_{\mathcal{O} \in \phi \times \phi} \lambda_{\mathcal{O}} C_I(x, P) \mathcal{O}^I(0) \quad (\text{OPE})$$

- ▶ Sum runs over *primary* \mathcal{O} 's
- ▶ $C_I(x, P)$ fixed by conformal symmetry [Dolan, Osborn '00]
- ▶ $\mathcal{O}^I = \mathcal{O}^{a_1 \dots a_l}$ can be any spin- l Lorentz representation (traceless symmetric tensor) with $l = 0, 2, \dots$
- ▶ Unitarity tells us that $\Delta_{\mathcal{O}} \geq l + 2 - \delta_{l,0}$ and that $\lambda_{\mathcal{O}}$ is real

CFT Review: Conformal Block Decomposition

Use OPE to evaluate 4-point function

$$\begin{aligned}
 & \langle \overbrace{\phi(x_1)\phi(x_2)} \overbrace{\phi(x_3)\phi(x_4)} \rangle \\
 &= \sum_{\mathcal{O} \in \phi \times \phi} \lambda_{\mathcal{O}}^2 C_I(x_{12}, \partial_2) C_J(x_{34}, \partial_4) \langle \mathcal{O}^I(x_2) \mathcal{O}^J(x_4) \rangle \\
 &\equiv \frac{1}{x_{12}^{2d} x_{34}^{2d}} \sum_{\mathcal{O} \in \phi \times \phi} \lambda_{\mathcal{O}}^2 g_{\Delta, l}(u, v)
 \end{aligned}$$

- ▶ $u = \frac{x_{12}^2 x_{34}^2}{x_{13}^2 x_{24}^2}$, $v = \frac{x_{14}^2 x_{23}^2}{x_{13}^2 x_{24}^2}$ conformally-invariant cross ratios.
- ▶ $g_{\Delta, l}(u, v)$ conformal block ($\Delta = \dim \mathcal{O}$ and $l = \text{spin of } \mathcal{O}$)

CFT Review: Conformal Blocks

Explicit formula [\[Dolan, Osborn '00\]](#)

$$g_{\Delta,l}(u,v) = \frac{(-1)^l}{2^l} \frac{z\bar{z}}{z-\bar{z}} [k_{\Delta+l}(z)k_{\Delta-l-2}(\bar{z}) - z \leftrightarrow \bar{z}]$$
$$k_{\beta}(x) = x^{\beta/2} {}_2F_1(\beta/2, \beta/2, \beta; x),$$

where $u = z\bar{z}$ and $v = (1-z)(1-\bar{z})$.

- ▶ Similar expressions in other even dimensions, recursion relations known in odd dimensions
- ▶ Alternatively can be viewed as eigenfunctions of the quadratic casimir of the conformal group [\[Dolan, Osborn '03\]](#)

CFT Review: Crossing Relations

- ▶ Four-point function $\langle \phi(x_1)\phi(x_2)\phi(x_3)\phi(x_4) \rangle$ is clearly symmetric under permutations of x_i
- ▶ After OPE, symmetry is non-manifest!
- ▶ Switching $x_1 \leftrightarrow x_3$ gives the “crossing relation”:

$$\sum_{\mathcal{O} \in \phi \times \phi} \lambda_{\mathcal{O}g\Delta,l}^2(u, v) = \left(\frac{u}{v}\right)^d \sum_{\mathcal{O} \in \phi \times \phi} \lambda_{\mathcal{O}g\Delta,l}^2(v, u)$$

$$\sum \text{Diagram 1} = \sum \text{Diagram 2}$$

- ▶ Other permutations give no new information

Outline

- ① CFT Review
- ② Bounds from Crossing Relations
- ③ Superconformal Blocks
- ④ Bounds on CFTs and SCFTs
- ⑤ Outlook

Review: Method of Rattazzi et. al. [arXiv:0807.0004]

- ▶ Let's study the OPE coefficient of a particular $\mathcal{O}_0 \in \phi \times \phi$
- ▶ We can rewrite crossing relation as

$$\underbrace{\lambda_{\mathcal{O}_0}^2 F_{\Delta_0, l_0}(u, v)}_{\mathcal{O}_0} = \underbrace{1}_{\text{unit op.}} - \underbrace{\sum_{\mathcal{O} \neq \mathcal{O}_0} \lambda_{\mathcal{O}}^2 F_{\Delta, l}(u, v)}_{\text{everything else}},$$

where

$$F_{\Delta, l}(u, v) \equiv \frac{v^d g_{\Delta, l}(u, v) - u^d g_{\Delta, l}(v, u)}{u^d - v^d}.$$

Review: Method of Rattazzi et. al. [arXiv:0807.0004]

Idea: Find a linear functional α such that

$$\begin{aligned}\alpha(F_{\Delta_0, l_0}) &= 1, \quad \text{and} \\ \alpha(F_{\Delta, l}) &\geq 0, \quad \text{for all other } \mathcal{O} \in \phi \times \phi.\end{aligned}$$

Applying to both sides:

$$\begin{aligned}\alpha(\lambda_{\mathcal{O}_0}^2 F_{\Delta_0, l_0}) &= \alpha\left(1 - \sum_{\mathcal{O} \neq \mathcal{O}_0} \lambda_{\mathcal{O}}^2 F_{\Delta, l}\right) \\ \lambda_{\mathcal{O}_0}^2 &= \alpha(1) - \sum_{\mathcal{O} \neq \mathcal{O}_0} \lambda_{\mathcal{O}}^2 \alpha(F_{\Delta, l}) \leq \alpha(1)\end{aligned}$$

since $\lambda_{\mathcal{O}}^2 \geq 0$ by unitarity.

Review: Method of Rattazzi et. al. [arXiv:0807.0004]

- ▶ To make the bound $\lambda_{\mathcal{O}_0}^2 \leq \alpha(1)$ as strong as possible:
Minimize $\alpha(1)$ subject to $\alpha(F_{\Delta_0, l_0}) = 1$ and $\alpha(F_{\Delta, l}) \geq 0$
- ▶ This is an infinite dimensional *linear programming problem...*
 to use known algorithms we must make it finite
- ▶ Can take α to be linear combinations of derivatives at some point in z, \bar{z} space


$$\alpha : F_{\Delta, l}(z, \bar{z}) \mapsto \sum_{m+n \leq N} a_{mn} \partial_z^m \partial_{\bar{z}}^n F_{\Delta, l}(1/2, 1/2)$$

- ▶ Discretize constraints to $\alpha(F_{\Delta_i, l_i}) \geq 0$ for $D = \{(\Delta_i, l_i)\}$
- ▶ Take $N \rightarrow \infty$ to recover “optimal” bound

Review: Method of Rattazzi et. al. [arXiv:0807.0004]

- ▶ Can make any assumptions about the spectrum that we want!
- ▶ E.g., can assume that all scalars appearing in the OPE $\phi \times \phi$ have dimension larger than some $\Delta_{\min} = \dim \mathcal{O}_0$
- ▶ If $\lambda_{\mathcal{O}_0}^2 \leq \alpha(1) < 0$, there is a contradiction with unitarity and the assumed spectrum can be ruled out

By scanning over different Δ_{\min} , one can obtain bounds on $\dim \phi^2$ as a function of $d = \dim \phi$

Bounds on $\dim \phi^2$ (taken from arXiv:0905.2211)

Limitations

- ▶ Single real ϕ , can't distinguish between \mathcal{O} 's with different global symmetry charges
- ▶ Example: chiral operator Φ in an $\mathcal{N} = 1$ SCFT
 - ▶ $\text{Re}[\Phi] \times \text{Re}[\Phi]$ contains \mathcal{O} 's from both $\Phi \times \Phi$ and $\Phi^\dagger \times \Phi$
 - ▶ $\Phi \times \Phi = \Phi^2 + \dots$, with $\dim \Phi^2 = 2 \dim \Phi$:
 Φ^2 satisfies bound and we learn nothing about $\Phi^\dagger \Phi$
- ▶ Supersymmetry also relates different conformal primaries, so we should additionally take this information into account

Let's try to generalize the method to deal with this case!

(see [Rattazzi, Rychkov, Vichi \[arXiv:1009.5985\]](#) for more on global symmetries)

Outline

- ① CFT Review
- ② Bounds from Crossing Relations
- ③ Superconformal Blocks**
- ④ Bounds on CFTs and SCFTs
- ⑤ Outlook

$\mathcal{N} = 1$ Superconformal Algebra

dim					
+1			P_a		
+1/2		Q_α		$\bar{Q}_{\dot{\alpha}}$	
0	$M_{\alpha\beta}$		D, R		$M_{\dot{\alpha}\dot{\beta}}$
-1/2		S_α		$\bar{S}_{\dot{\alpha}}$	
-1			K_a		

$$\{Q, \bar{Q}\} = P$$

$$\{S, \bar{S}\} = K$$

- ▶ Superconformal primary means $[S, \mathcal{O}(0)] = [\bar{S}, \mathcal{O}(0)] = 0$
- ▶ Descendants obtained by acting with P, Q, \bar{Q}
- ▶ Chiral means $[\bar{Q}, \phi(0)] = 0$

Superconformal Block Decomposition

ϕ : scalar chiral superconformal primary of dimension d in an SCFT
(lowest component of chiral superfield Φ)

$$\langle \overbrace{\phi(x_1)\phi^\dagger(x_2)} \overbrace{\phi(x_3)\phi^\dagger(x_4)} \rangle = \frac{1}{x_{12}^{2d}x_{34}^{2d}} \sum_{\mathcal{O} \in \Phi \times \Phi^\dagger} |\lambda_{\mathcal{O}}|^2 (-1)^l \mathcal{G}_{\Delta, l}(u, v)$$

- ▶ Sum over superconformal primaries \mathcal{O}^I with zero R -charge
- ▶ $\lambda_{\mathcal{O}}$ real for even spin \mathcal{O}^I , imaginary for odd spin \mathcal{O}^I
- ▶ $x_1 \leftrightarrow x_3$ gives crossing relation only involving $\mathcal{O}^I \in \Phi \times \Phi^\dagger$
- ▶ Must organize superconformal descendants into reps of the conformal subalgebra...

Superconformal Block Derivation

Multiplet built from \mathcal{O} (generically) contains four conformal primaries with vanishing R -charge and definite spin:

name	operator	dim	spin
\mathcal{O}	\mathcal{O}	Δ	l
J, N	$Q\bar{Q}\mathcal{O} + \#P\mathcal{O}$	$\Delta + 1$	$l + 1, l - 1$
D	$Q^2\bar{Q}^2\mathcal{O} + \#PQ\bar{Q}\mathcal{O} + \#PP\mathcal{O}$	$\Delta + 2$	l

- ▶ Superconformal symmetry fixes coefficients of $\langle\phi\phi^\dagger J\rangle, \langle\phi\phi^\dagger N\rangle, \langle\phi\phi^\dagger D\rangle$ in terms of $\langle\phi\phi^\dagger \mathcal{O}\rangle$
- ▶ Must also normalize J, N, D to have canonical 2-pt functions
- ▶ Superconformal block is then a sum of $g_{\Delta, l}$'s for \mathcal{O}, J, N, D

Superconformal Block Derivation

We find,¹

$$\mathcal{G}_{\Delta,l} = g_{\Delta,l} - \frac{(\Delta + l)}{2(\Delta + l + 1)}g_{\Delta+1,l+1} - \frac{(\Delta - l - 2)}{8(\Delta - l - 1)}g_{\Delta+1,l-1} + \frac{(\Delta + l)(\Delta - l - 2)}{16(\Delta + l + 1)(\Delta - l - 1)}g_{\Delta+2,l}$$

- ▶ When unitarity bound $\Delta \geq l + 2$ is saturated, multiplet is shortened
- ▶ $\mathcal{G}_{\Delta,l}$ can also be determined from consistency with $\mathcal{N} = 2$ superconformal blocks computed by [\[Dolan, Osborn '01\]](#)

¹after plenty of algebra

Outline

- ① CFT Review
- ② Bounds from Crossing Relations
- ③ Superconformal Blocks
- ④ Bounds on CFTs and SCFTs
- ⑤ Outlook

Bounds on Dimension of $\Phi^\dagger\Phi$

Isolating the lowest dimension scalar $\Phi^\dagger\Phi \in \Phi \times \Phi^\dagger$, we have

$$|\lambda_{\Phi^\dagger\Phi}|^2 \mathcal{F}_{\Delta_{\min},0} = 1 - \sum_{\mathcal{O} \neq \Phi^\dagger\Phi} |\lambda_{\mathcal{O}}|^2 \mathcal{F}_{\Delta,l},$$


where $\Delta_{\min} = \dim \Phi^\dagger\Phi$, and $\mathcal{F}_{\Delta,l}$ is $F_{\Delta,l}$ with $g_{\Delta,l} \rightarrow (-1)^l \mathcal{G}_{\Delta,l}$.

Now minimize $\alpha(1)$ subject to

- ▶ $\alpha(\mathcal{F}_{\Delta,0}) \geq 0$ for all $\Delta \geq \Delta_{\min}$,
- ▶ $\alpha(\mathcal{F}_{\Delta,l}) \geq 0$ for all $\Delta \geq l + 2$ and $l \geq 1$,
- ▶ $\alpha(\mathcal{F}_{\Delta_{\min},0}) = 1$

If $\alpha(1) < 0$, we get $|\lambda_{\Phi^\dagger\Phi}|^2 < 0 \implies \Phi^\dagger\Phi$ can't have $\dim \Delta_{\min}$

Upper Bound on Dimension of $\Phi^\dagger\Phi$



- ▶ Scanning over Δ_{\min} , minimizing $\alpha(1)$ over 21 dimensional space of derivatives

Flavor Currents

- ▶ If ϕ transforms under flavor symmetry with charges T^I , conserved currents J^I appear in the $\phi \times \phi^\dagger$ OPE:

$$\langle \phi \phi^\dagger J^I \rangle \sim -\frac{i}{2\pi^2} T^I \quad (\text{Ward id.})$$



- ▶ Flavor current conformal blocks are then determined by current 2-pt functions

$$\langle J^I J^J \rangle \sim \frac{3}{4\pi^4} \tau^{IJ}$$

$$\langle \phi \phi^\dagger \phi \phi^\dagger \rangle \sim -\frac{1}{3} \tau_{IJ} T^I T^J g_{3,1} \quad (\text{general CFTs}),$$

$$\langle \phi \phi^\dagger \phi \phi^\dagger \rangle \sim \tau_{IJ} T^I T^J \mathcal{G}_{2,0} \quad (\text{SCFTs}),$$

where $\tau_{IJ} = (\tau^{IJ})^{-1}$ (in SCFTs, $\tau^{IJ} = -3\text{Tr}(RT^I T^J)$).

Upper Bounds on $\tau_{IJ}T^IT^J$ 

- ▶ Example: SUSY QCD with $\frac{3}{2}N_c < N_f < 3N_c$, $M = Q\tilde{Q}$
 $\langle MM^\dagger MM^\dagger \rangle$: $d = 3 - \frac{3N_c}{N_f}$ and $\tau_{IJ}T^IT^J = \frac{2}{3} \frac{N_f - 1}{N_c^2}$
- ▶ In dual AdS_5 , $(8\pi^2 L)\tau_{IJ} = g_{IJ}^2$. Gauge coupling can't be too strong in presence of charged scalar.



The Stress Tensor

- ▶ Ward identity ensures $T^{ab} \in \phi \times \phi$
- ▶ $\langle TT \rangle$ is proportional to the central charge c
(trace anomaly $16\pi^2 \langle T_a^a \rangle = c(\text{Weyl})^2 - a(\text{Euler})$)
- ▶ In an SCFT, T lives in the supercurrent multiplet
 $\mathcal{J}^a = J_R^a + \theta \sigma_b \bar{\theta} T^{ab} + \dots$, and c determined by $U(1)_R$
- ▶ Conformal block contributions are

$$\langle \phi \phi \phi \phi \rangle \sim \frac{d^2}{90c} g_{4,2} \quad (\text{general CFTs})$$

$$\langle \phi \phi^\dagger \phi \phi^\dagger \rangle \sim -\frac{d^2}{36c} \mathcal{G}_{3,1} \quad (\text{SCFTs})$$

Lower Bound on c in General CFT



- ▶ In dual AdS_5 , $c \sim \pi^2 L^3 M_P^3$. Gravity can't be arbitrarily strong in presence of light bulk scalar!

(See also [Rattazzi, Rychkov, Vichi \[arXiv:1009.2725\]](#))

Outline

- ① CFT Review
- ② Bounds from Crossing Relations
- ③ Superconformal Blocks
- ④ Bounds on CFTs and SCFTs
- ⑤ Outlook

Outlook

We calculated:

- ▶ Superconformal blocks
- ▶ Bound $\dim \Phi^\dagger \Phi < f_{\Phi^\dagger \Phi}(d)$
- ▶ Bound $\tau_{IJ} T^I T^J \leq f_\tau(d)$ in CFT, SCFT
- ▶ Bound $c \geq f_c(d)$ in CFT, SCFT

In the future, we'd like:

- ▶ Stronger bounds to make contact with BSM motivation!
- ▶ Better algorithms (esp. to deal with global symmetries)
- ▶ SUSY theories that come close to saturating bounds on τ, c .
- ▶ Bounds in other numbers of dimensions.
- ▶ Understand bounds from bulk dual perspective.