

# Bottom-up naturalness

Sonia El Hedri

with


Anson Hook


November 20, 2013

arXiv:1305.6608

# The hierarchy problem

- ▶ The Higgs mass in the SM is not protected by symmetries
- ▶ One loop contributions quadratically divergent (top, gauge)
- ▶ Mass corrections of order the cutoff scale  $\Lambda^2$
- ▶ New physics at the TeV scale


$$\propto -m_t^2 \Lambda^2$$


$$\propto +m_V^2 \Lambda^2$$

# Traditional approaches


## Top down approaches

Assuming a high energy mechanism which cancels the divergences  
at all loop levels


- ▶ SUSY
- ▶ Extra dimensions
- ▶ Little Higgs
- ▶ ...

# Traditional approaches

New particles running in loops




Cancellation terms give new signatures



# Traditional approaches

But dominant signatures from other terms



Model dependent

Not directly related to the quadratic divergences

# Bottom up approach?

Study low energy signatures of naturalness

- ⇒ Cancellation at one loop only
- ⇒ No complete model

But


- ⇒ Necessary conditions for naturalness
- ⇒ Model independent approach
- ⇒ Hints for new complete theories?
- ⇒ Limited number of simplified models

# Minimal naturalness

Naturalness is enforced by

$$y' H \psi_1 \psi_2$$

$$\lambda H^\dagger H \psi^\dagger \psi$$




- ▶ Find all possible  $\psi$
- ▶ For each  $\psi$ , look for signatures which vanish when  $y$  or  $\lambda$  vanishes

# The trilinear term

# Properties of $\psi_1, \psi_2$

$$\mathcal{L} = yH\psi_1\psi_2$$




- ▶  $\psi_1$  and  $\psi_2$  are fermions
- ▶ Negative one loop contribution
- ▶  $\psi_1$  and/or  $\psi_2$  charged under **at least**  $SU(2)$

# Trilinear term – phenomenology

$\psi_1$  and  $\psi_2$  non SM


- ▶ Electroweakino-like phenomenology



- ▶ Decays to Higgs and gauge bosons
- ▶  $\cancel{E}_T$ , CHAMPs, R-hadrons...

$\psi_1$  SM

- ▶ Fourth generation or RH neutrino



- ▶ Pair or single production
- ▶ Decays to  $W$ ,  $Z$ ,  $h$  + SM partner

## The quartic term

$$\mathcal{L} = \mathcal{L}_{\text{SM}} + \lambda H^\dagger H \psi^\dagger \psi$$


# The quartic term

$$\mathcal{L}_2 = \lambda H^\dagger H \psi^\dagger \psi$$

$$\supset \lambda v h \psi^\dagger \psi + \frac{\lambda}{2} h^2 \psi^\dagger \psi$$



Scalar



Vector-like fermion

- ▶ New Higgs decay modes
- ▶  $\psi$  is a dark matter particle
- ▶  $\psi$  gets a vev
- ▶  $\psi$  is charged under the SM


# The quartic term

$$\mathcal{L}_2 = \lambda H^\dagger H \psi^\dagger \psi$$

$$\supset \lambda v h \psi^\dagger \psi + \frac{\lambda}{2} h^2 \psi^\dagger \psi$$



Scalar





Vector-like fermion

- ▶ New Higgs decay modes
- ▶  $\psi$  is a dark matter particle
- ▶  $\psi$  gets a vev
- ▶  $\psi$  is charged under the SM

# Higgs decays to $\psi^\dagger \psi$

ATLAS-CONF-2013-011



- ▶ Invisible decay modes
- ▶ Top and gauge divergences  $\Rightarrow$  Excluded
- ▶ Other divergences  $\Rightarrow$  Effect too small

# The quartic term

$$\mathcal{L}_2 = \lambda H^\dagger H \psi^\dagger \psi$$

$$\supset \lambda v h \psi^\dagger \psi + \frac{\lambda}{2} h^2 \psi^\dagger \psi$$



Scalar




Vector-like fermion


- ▶ ~~New Higgs decay modes~~
- ▶  $\psi$  is a dark matter particle
- ▶  $\psi$  gets a vev
- ▶  $\psi$  is charged under the SM

$\psi$  is dark matter

Direct detection



Annihilation



Spin independent interactions


Higgs portal only

$\psi\psi \rightarrow hh, WW, ZZ$

$$\lambda_f = \frac{N_c y^2}{2 N_f m_\psi}$$


$$\lambda_s = \frac{2 N_c}{N_s} y^2$$

# Relic abundance




Non thermal production or subdominant DM component

# Indirect detection



- ▶ No top quadratic divergences cancellation
- ▶ Gauge cancellation possible for  $m_\psi > 500$  GeV

# Direct detection



- ▶ Top and gauge cancellation excluded

# $\psi$ dark matter

Correlated direct and indirect detection signatures

- ▶ If fermion, direct detection signature but no indirect detection signal
- ▶ If scalar,

$$\frac{\sigma_{\text{SI}}}{\langle \sigma v \rangle_{v=0}} = \frac{16f^2 m_p^2}{m_h^4} = 1.5 \times 10^{-19} \frac{\text{cm}^2}{\text{cm}^3/\text{s}}$$

Measurable at FERMI, XENON100, LUX


Sub-TeV  $\psi$  cannot cancel the top quadratic divergences

Small region still left for gauge quadratic divergences


# The quartic term

$$\mathcal{L}_2 = \lambda H^\dagger H \psi^\dagger \psi$$

$$\supset \lambda v h \psi^\dagger \psi + \frac{\lambda}{2} h^2 \psi^\dagger \psi$$



Scalar



Vector-like fermion

- ▶ ~~New Higgs decay modes~~
- ▶  ~~$\psi$  is a dark matter particle~~
- ▶  $\psi$  gets a vev
- ▶  $\psi$  is charged under the SM

# Scalar with a vev

$$\mathcal{L} = \lambda vv_\psi h\psi + \frac{\lambda}{2}v_\psi\psi hh + \frac{\lambda}{2}vh\psi\psi + \dots$$

If  $\psi$  is an  $SU(2)$  doublet  $\Rightarrow$  two Higgs doublet model

What about a singlet?

- ▶  $h$  decays (already studied)
- ▶ Mixing with the Higgs  $\Rightarrow \cos \alpha > 0.93$
- ▶  $\psi$  decays  $\Rightarrow \sin \alpha < 0.20$
- ▶ For our minimal model, top cancellation requires

$$v_\psi > 2 \text{ TeV}$$


# The quartic term

$$\mathcal{L}_2 = \lambda H^\dagger H \psi^\dagger \psi$$

$$\supset \lambda v h \psi^\dagger \psi + \frac{\lambda}{2} h^2 \psi^\dagger \psi$$



Scalar




Vector-like fermion

- ▶ ~~New Higgs decay modes~~
- ▶  ~~$\psi$  is a dark matter particle~~
- ▶  ~~$\psi$  gets a vev~~
- ▶  $\psi$  is charged under the SM

# $\psi$ charged under the SM

$$\mathcal{L} \supset \lambda h \psi^\dagger \psi + g_G V_G^\mu \gamma_\mu \psi^\dagger \psi$$



- ▶ One loop Higgs couplings to gauge bosons modified

$SU(3)$  production, not visible

$SU(2)$  decay, hard to reach at the LHC

$U(1)_{\text{EM}}$  decay, high luminosity LHC


## Example: electrically charged $\psi$

$\psi$  has electric charge  $Q$  and cancels the top quadratic divergences

$$\mathcal{L} \supset -m\psi^\dagger\psi + \frac{3y_t^2}{2m}\psi^\dagger\psi hh$$

$$\frac{\Gamma(h \rightarrow \gamma\gamma)}{\Gamma_{\text{SM}}(h \rightarrow \gamma\gamma)} = \left| 1 + \frac{Q^2}{6.49} \frac{4}{3} \frac{\partial \log m_\psi}{\partial \log v} \left( 1 + \frac{7m_h^2}{120m_\psi^2} \right) \right|^2$$

## Example: electrically charged $\psi$



Less than 10% modifications at high mass

# Quartic term: summary

$$\lambda \psi^\dagger \psi H^\dagger H \supset \begin{cases} \lambda v h \psi^\dagger \psi + \frac{\lambda}{2} h h \psi^\dagger \psi \\ \frac{\lambda}{2} v v_\psi h \psi + \lambda v h \psi^\dagger \psi + \lambda v_\psi \psi h h \end{cases}$$

- ▶  $\psi$  light
  - ▶ Invisible Higgs decays
  - ▶ Cannot cancel top and gauge quadratic divergences
- ▶  $\psi$  dark matter
  - ▶ Correlated direct and indirect detection signatures
  - ▶ Strong constraints on top and gauge divergences cancellation
- ▶  $\psi$  scalar with a vev
  - ▶ Precision Higgs coupling measurements
  - ▶ Tight constraints on  $v_\psi$
- ▶  $\psi$  charged under the SM
  - ▶ One loop contributions to  $h \rightarrow VV$
  - ▶ Modifications too small to observe with current searches

# Current prospects

- ▶ Strong constraints in specific cases for top and gauge cancellation (dark matter, light particle, etc...)
- ▶ In most cases, precision Higgs measurements are needed

Most minimal signatures cannot be observed with current experiments!

Can some simple extensions be probed at the LHC?

# Finding minimal extensions

Minimal naturalness – Quartic term extension

$$\mathcal{L} = \mathcal{L}_{\text{SM}} + \lambda H^\dagger H \psi^\dagger \psi$$


Find additional terms:

- ▶ IR effect
- ▶ No assumptions about the UV physics
- ▶ New decay modes for  $\psi$ , new LHC signatures
- ▶ **Signatures vanish when  $\lambda \rightarrow 0$**

# Mass mixing

Only possible term

$$\mathcal{L} \supset m_0 \psi^\dagger \psi_{\text{NEW}}$$



$\psi_{\text{NEW}}$  {

- Higgs boson  $\Rightarrow$  2HDM
- Stable new particle  $\Rightarrow \cancel{\not{E}_T}$ , CHAMPs, R-hadrons
- SM fermion

# Mass mixing

$$\mathcal{L} \supset \lambda_1 \psi^\dagger \psi H^\dagger H + \lambda_2 \psi_{\text{NEW}}^\dagger \psi_{\text{NEW}} H^\dagger H$$

- ▶ Measuring  $\lambda_2$   
⇒ **Indirect** evidence of  $\lambda_1$
- ▶ Three-body decays to  $\psi_{\text{NEW}}$ ,  $WW$ ,  $hh$  and  $ZZ$
- ▶ Two-body decay to  $\psi_{\text{NEW}}$  and  $h$
- ▶ NO two-body decays to gauge bosons

# Example: Little Higgs model

Fermionic top partner

$$\mathcal{L} = m_\psi \psi \psi^c + \lambda_1 \psi^c H Q + \lambda_2 u^c H Q + \frac{\lambda_3}{m_\psi} \psi \psi^c H^\dagger H$$

In mass basis

$$\begin{aligned} \mathcal{L} = & m_T T T^c + \lambda_T T^c H Q + y_t u^c H Q \\ & + \frac{\lambda_{TT}}{m_T} T^c T H^\dagger H + \frac{\lambda_{tT}}{m_T} u^c T H^\dagger H \end{aligned}$$

- ▶  $\lambda_{tT}$  generated by  $\lambda_{TT}$  and mass mixing

## Example: Little Higgs model


$$\mathcal{L} \supset \lambda_T T^c H Q + \frac{\lambda_{tT}}{m_T} u^c T H^\dagger H$$

- ▶ Two-body decays from trilinear + quartic terms
- ▶  $\lambda_T$  usually expected to dominate
- ▶ But two-body signatures dominantly from quartic if

$$\lambda_{tT} > \lambda_T \frac{m_T}{v}$$

Little Higgs is a good example for large quartic and moderate  $m_T$

# $T$ decay modes



$T \rightarrow tH$  largely dominating

# Vector-like fermions at the LHC

$$\mathcal{L} \supset \lambda \psi \psi_{\text{NEW}} H^\dagger H \supset \lambda_1 \psi \psi_{\text{NEW}} h$$

- ▶ Choose  $\psi_{\text{NEW}}$  SM fermion
- ▶ Consider only two-body decays
- ▶ Derive bounds for top quark, light quark and lepton partners


$$T \rightarrow t + h$$

$$U \rightarrow u + h$$

$$L \rightarrow l + h$$

# Top quark partners

- ▶ ATLAS-CONF-2013-018
- ▶ 8 TeV,  $14.3\text{fb}^{-1}$



- ▶  $\text{Br}(T \rightarrow th) = 100\% \Rightarrow m_T > 850\text{ GeV}$

# Light quark partner


- ▶  $hhjj$  final state
- ▶ No rapidity gap between the two jets
- ▶ Low branching ratio to leptons + low lepton ID efficiency
- ▶  $h \rightarrow \gamma\gamma$  search does not veto on extra jets
- ▶ Current bounds on  $\gamma\gamma$  allow up to 10 pb signal

$$m_U > 300 \text{ GeV}$$

# Lepton partner

- ▶  $l^+ l^- hh$  final state
- ▶  $l^+ l^- + 2b$  dominant but existing searches require an on-shell  $Z$
- ▶ 4-lepton events from  $h \rightarrow W^+ W^-, \tau\tau$
- ▶ ATLAS-CONF-2013-036
- ▶ 4-leptons + effective mass cut
- ▶ Low background, high signal efficiency

# Lepton partner




Low production cross section  $\Rightarrow$  no exclusion bounds

# Summary

- ▶ Two possible operators to cancel one-loop divergences
- ▶ Bottom-up approach: study signatures which vanish when these operators vanish
- ▶ New Yukawa term  $\Rightarrow$  electroweakino phenomenology, CHAMPs, R-hadrons
- ▶ Quartic term
  - ▶ Correlated dark matter detection signatures
  - ▶ Higgs precision measurements
- ▶ Mass mixing with a SM fermion gives new decay modes
- ▶ Only one two-body decay mode to SM fermion + Higgs
- ▶ Strong bounds on top partners at the LHC, high luminosity + dedicated searches needed for the other particles

# Goldstone boson equivalence theorem



$$|M(T \rightarrow thh)|^2 \sim \frac{\lambda^2}{2} p_{T,\mu} p_t^\mu$$

$$|M(T \rightarrow tW^+W^-)|^2 \sim 4\lambda^2 m_W^4 p_{T,\mu} p_t^\mu \frac{1}{((p_T - p_t)^2 - m_h^2)^2} \frac{(p_{W^+} \cdot p_{W^-})^2}{m_W^4}$$

# Little Higgs model

$$\Sigma = \exp \left( \frac{i}{f} \begin{pmatrix} 0 & H \\ H^\dagger & 0 \end{pmatrix} \right) \begin{pmatrix} 0 \\ f \end{pmatrix}$$

After symmetry breaking

$$\mathcal{L} \supset \lambda_1 u_3^c \Sigma \chi + \lambda_2 f u'^c u'$$

At lowest order

$$\mathcal{L} \supset f(\lambda_1 u_3^3 + \lambda_2 u'^c) u' - \lambda_1 u_3^c H Q_3 + \frac{\lambda_1}{2f} H H^\dagger u_3^c u'$$

# Little Higgs model

After diagonalization

$$\mathcal{L} \supset \frac{\lambda_1 \lambda_2}{\sqrt{\lambda_1^2 + \lambda_2^2}} t_3^c H Q_3 + \frac{\lambda_1^2}{\sqrt{\lambda_1^2 + \lambda_2^2}} T^c H Q_3 + \frac{\lambda_1^2}{2m_T} H H^\dagger T^c T + \frac{\lambda_1 \lambda_2}{2m_T} H H^\dagger t_3^c T$$

Two body decays from the quartic term dominate if

$$\frac{\lambda_2^2 v}{\sqrt{2} y m_T} \gg 1$$