Gauge Mediation from Emergent SUSY

Siew-Phang Ng
Bartol Research Institute.


[Goh, SPN & Okada(hep-ph/0511301)]
[Goh, SPN & Okada(in progress)]

---

*Presented at the Santa Fe Summer Workshop on Particle Theory and the LHC, Santa Fe, New Mexico on July 23-29, 2006.

Introduction

LHC I: Overview

- Turning on in 2007.

LHC I: Overview

- Turning on in 2007.
- Herald of a new dawn? Or something completely different...

LHC II: Operation

- Typically in 1 second, we get 10^9 events. Applying the triggers...

LHC II: Operation

- Typically in 1 second, we get 10^9 events. Applying the triggers...
  - Level 1: $\sim 10^7$ events
  - Level 2: $\sim 10^5$ events
  - Level 3: $\sim 10^2$ events

LHC II: Operation

- Typically in 1 second, we get 10^9 events. Applying the triggers...
  - Level 1: $\sim 10^7$ events
  - Level 2: $\sim 10^5$ events
  - Level 3: $\sim 10^2$ events (recorded)

LHC II: Operation

- Typically in 1 second, we get 10^9 events. Applying the triggers...
  - Level 1: $\sim 10^7$ events
  - Level 2: $\sim 10^5$ events
  - Level 3: $\sim 10^2$ events (recorded)
- Very real danger of, as it were, "throwing out the baby with the bathwater".

- Incidentally, no babies were harmed in the making of this presentation.

Path Ahead

- What can we do about it?
  - Hidden valley (Zurek, Strassler)
  - Exotic Higgs decay (Fox, Chang, Weiner)
  - etc.

Path Ahead

- What can we do about it?
  - Hidden valley (Zurek, Strassler)
  - Exotic Higgs decay (Fox, Chang, Weiner)
  - etc.
- In short, more than ever, there's a need to question conventional wisdom.
- So what am I trying to sell?

Path Ahead

- What can we do about it?
  - Hidden valley (Zurek, Strassler)
  - Exotic Higgs decay (Fox, Chang, Weiner)
  - etc.
- In short, more than ever, there's a need to question conventional wisdom.
- So what am I trying to sell?Gauge mediation from emergent susy, of course. A new approach to susy model building.

Typical Gauge Mediation

- Setup and properties
  - Parameters: F, M and $\Lambda = F/M$
  - Dynamical Susy Breaking
  - Phenomenological viability constrains F, M and $\Lambda = F/M$
  - Gravitino LSP
  - Bounds from cosmology
  - Problems with DSB.
- How is GMES different?

Big Picture

Big Picture I: GMES

- Based on the paradigm of Susy without Susy. [Goh, Luty & SPN(hep-th/0309103)]

Big Picture I: GMES

- Based on the paradigm of Susy without Susy. [Goh, Luty & SPN(hep-th/0309103)]
- Susy is an ACCIDENTAL symmetry! (More on this later)

Big Picture I: GMES

- Based on the paradigm of Susy without Susy. [Goh, Luty & SPN(hep-th/0309103)]
- Susy is an ACCIDENTAL symmetry! (More on this later)
- Original realization does not solve the Susy flavor problem.

Big Picture I: GMES

- Based on the paradigm of Susy without Susy. [Goh, Luty & SPN(hep-th/0309103)]
- Susy is an ACCIDENTAL symmetry! (More on this later)
- Original realization does not solve the Susy flavor problem.
- So we decided to use gauge mediation.

Big Picture II: CFT

- Key assumption: We live in a superconformal basin.

Big Picture II: CFT

- Key assumption: We live in a superconformal basin.



- Features
  - Start at the edge

Big Picture II: CFT

- Key assumption: We live in a superconformal basin.



- Features
  - Start at the edge
  - Flows towards the fixed point

Big Picture II: CFT

- Key assumption: We live in a superconformal basin.



- Features
  - Start at the edge
  - Flows towards the fixed point
  - Flow terminated before f.p.

Big Picture II: CFT

- Key assumption: We live in a superconformal basin.



- Features
  - Start at the edge
  - Flows towards the fixed point
  - Flow terminated before f.p.
- Subtleties
  - Susy breaking operators
  - Anomalous dimensions
  - Fundamental vs Emergent fields
  - Emergent Susy

Big Picture II: CFT

- Key assumption: We live in a superconformal basin.



- Features
  - Start at the edge
  - Flows towards the fixed point
  - Flow terminated before f.p.
- Subtleties
  - Susy breaking operators
  - Anomalous dimensions
  - Fundamental vs Emergent fields
  - Emergent Susy

- For more insights, use AdS-CFT dictionary.

Big Picture III: AdS dual

- Key assumption: There are no light bulk scalars.

Big Picture III: AdS dual

- Key assumption: There are no light bulk scalars.



- Features
  - UV brane

Big Picture III: AdS dual

- Key assumption: There are no light bulk scalars.



- Features
  - UV brane
  - Bulk

Big Picture III: AdS dual

- Key assumption: There are no light bulk scalars.



- Features
  - UV brane
  - Bulk
  - IR brane

Big Picture III: AdS dual

- Key assumption: There are no light bulk scalars.



- Features
  - UV brane
  - Bulk
  - IR brane
- Subtleties
  - Susy breaking transmission
  - Bulk scalar masses
  - Bulk vs IR-localized fields
  - Emergent Susy

Big Picture III: AdS dual

- Key assumption: There are no light bulk scalars.



- Features
  - UV brane
  - Bulk
  - IR brane
- Subtleties
  - Susy breaking transmission
  - Bulk scalar masses
  - Bulk vs IR-localized fields
  - Emergent Susy

- Much easier to construct an explicit example on the AdS side.

Setup

Setup I: Explicit Model

- Randall-Sundrum model on a $S^1/Z_2 \times Z_2$ orbifold

$$ds^2 = e^{-2\sigma(y)}\eta_{\mu\nu}dx^\mu dx^\nu + dy^2.$$

- Action is given by

$$S = \frac{M_5^3}{k}\int d^4x \int d^4\theta(\omega^\dagger\omega - \varphi^\dagger\varphi) + \int d^4x \int_0^\ell dy L_{hyp},$$

where $\omega = e^{-k\ell} + \cdots + \theta^2 F_\omega$ and $\varphi = 1 + \theta^2 F_\varphi$.

- Hypermultiplet action is

$$\begin{aligned} L_{hyp} \;=\; & \int d^4\theta e^{-2\sigma}(\Phi^\dagger\Phi + \tilde{\Phi}^\dagger\tilde{\Phi}) + \left[\int d^2\theta e^{-3\sigma}\left(\tfrac{1}{2}\tilde{\Phi}\overleftrightarrow{\partial}_y\Phi\right.\right. \\ & \left.\left. + c\sigma'\tilde{\Phi}\Phi\right) + \text{h.c.}\right] - \delta(y)U(\Phi,\tilde{\Phi},F,\tilde{F}) \\ & + \delta(y-\ell)\omega^3\left[\int d^4\theta\, W(\Phi,\tilde{\Phi}) + \text{h.c.}\right] \end{aligned}$$

Setup II: EoM Solutions

- General solution (for $0 < y < \ell$) is

$$
\begin{aligned}
F &= F_0 e^{-(c-\frac{3}{2})\sigma} \\
\tilde{F} &= \tilde{F}_0 \frac{\sigma'}{k} e^{(c+\frac{3}{2})\sigma} \\
\Phi &= \Phi_0 e^{-(c-\frac{3}{2})\sigma} - \frac{\tilde{F}_0^\dagger}{(2c+1)k} e^{(c+\frac{5}{2})\sigma} \\
\tilde{\Phi} &= \tilde{\Phi}_0 e^{(c+\frac{3}{2})\sigma} - \frac{F_0^\dagger}{(2c-1)k} e^{-(c-\frac{5}{2})\sigma}
\end{aligned}
$$

- The prefactors are determined from the junction conditions.
- Digression: AdS-CFT dictionary

$$\dim(O_{\Phi,\tilde{\Phi}}) = d, \tilde{d} = 2 + |c \pm \tfrac{1}{2}|$$

SUSY breaking

SUSY breaking I: Various Mechanisms

- 5-d gravity loop contribution is $m_{\text{gravity}} \sim \omega^2$. [Gregoire et al(hep-th/0411216)]
- Effective 4-d Lagrangian that characterizes the soft SUSY breaking masses from the various mechanisms

$$L_{soft} = -V_{\text{eff},\omega} + \int d^4\theta \omega^\dagger \omega \left[1 + (1 + \Phi_{\text{IR}}^\dagger \Phi_{IR}) (Q^\dagger Q + X^\dagger X + \bar{X}^\dagger \bar{X})\right] + \int d^2\theta\, \omega^3 \Phi \bar{X} X + \text{h.c.}$$

- For the models of interest, scale of anomaly mediation is $m_{\text{anomaly}} \sim \frac{F_\omega}{\omega} = \frac{1}{\omega}\frac{\partial V_{\text{eff},\omega}}{\partial \omega} \sim \Lambda_{\text{IR}} \omega^{d-5}$. [Luty & Sundrum(hep-th/0012158)]
- After canonical normalization, direct mediation contributes $m^2_{\text{direct}} \sim F^\dagger_{\text{IR}} F_{\text{IR}}$. Generally, flavor non-diagonal. [Goh, Luty & SPN(hep-th/0309103)]
- What about gauge mediation?

SUSY breaking II: Gauge Mediation

- Mass matrix of the scalar messengers is completely specified.

$$m^2_{\text{messenger}} = \begin{pmatrix} \omega^\dagger\omega|\Phi_{\text{IR}}|^2 + |F_{\text{IR}}|^2 & \omega F_{\text{IR}} \\ \omega^\dagger F^\dagger_{\text{IR}} & \omega^\dagger\omega|\Phi_{\text{IR}}|^2 + |F_{\text{IR}}|^2 \end{pmatrix}$$

- Scale of gauge mediation is $m_{\text{gauge}} \sim \frac{F_{\text{IR}}}{\Phi_{\text{IR}}}$ subject to certain constraints.
- For a particular class of theories, we have

$$m_{\text{soft}} \sim \begin{cases} \frac{F_{\text{IR}}}{\Phi_{\text{IR}}} & \sim \Lambda_{\text{IR}}\omega^{\frac{d-5}{3}} & gauge & \checkmark \Rightarrow d > 5 \\ F_{\text{IR}} & \sim \Lambda_{\text{IR}}\omega^{\frac{2(d-5)}{3}} & direct & subdom. \\ \frac{F_\omega}{\omega} & \sim \Lambda_{IR}\omega^{d-5} & anomaly & subdom. \\ & \sim \Lambda_{IR}\omega & gravity & subdom? \end{cases}$$

Stabilization

Stabilization I: Brane Potentials

- For gauge mediation to dominate, the following is required: $(+,+)$ orbifold parity, $d > 5$ ($c < -\frac{5}{2}$) and the potentials

$$U = b(\Phi_{\rm UV} + \Phi_{\rm UV}^\dagger), \qquad W = a\Phi_{\rm IR}^3$$

- Effective potential is

$$V_{\rm eff} = \frac{3b}{4}\Phi_{\rm IR}\omega^{d-1} + \ldots = -A\omega^{4\frac{d-2}{3}} + \ldots$$

where $A > 0$.

Stabilization I: Brane Potentials

- For gauge mediation to dominate, the following is required: $(+,+)$ orbifold parity, $d > 5$ ($c < -\frac{5}{2}$) and the potentials

$$U = b(\Phi_{\rm UV} + \Phi_{\rm UV}^\dagger), \qquad W = a\Phi_{\rm IR}^3$$

- Effective potential is

$$V_{\rm eff} = \frac{3b}{4}\Phi_{\rm IR}\omega^{d-1} + \ldots = -A\omega^{4\frac{d-2}{3}} + \ldots$$

where $A > 0$.

- Introduce Ψ :$(+,+)$ orbifold parity, $c > 0$ (good only for stabilization) and potentials

$$U = b'\Psi_{\rm UV}^2 + b_2' F + \text{h.c.}, \qquad W = a'\Psi_{\rm IR}^2$$

- Hence for stabilization, $d' \gtrsim \frac{2d+5}{3}$.
- Checked SUSY breaking.

Stabilization I: Brane Potentials

- For gauge mediation to dominate, the following is required: $(+,+)$ orbifold parity, $d > 5$ ($c < -\frac{5}{2}$) and the potentials

$$U = b(\Phi_{\rm UV} + \Phi_{\rm UV}^\dagger), \qquad W = a\Phi_{\rm IR}^3$$

- Effective potential is

$$V_{\rm eff} = \frac{3b}{4}\Phi_{\rm IR}\omega^{d-1} + \ldots = -A\omega^{4\frac{d-2}{3}} + \ldots$$

where $A > 0$.

- Introduce Ψ :$(+,+)$ orbifold parity, $c > 0$ (good only for stabilization) and potentials

$$U = b'\Psi_{\rm UV}^2 + b_2' F + \text{h.c.}, \qquad W = a'\Psi_{\rm IR}^2$$

- Hence for stabilization, $d' \gtrsim \frac{2d+5}{3}$.
- Checked SUSY breaking. ✓

Stabilization II: Phenomenology

- Combining FCNC and Casimir constraints, and taking $m_{\rm soft} \sim 100$ GeV and $M_5 = 2.4 \times 10^{18}$ GeV,

$$\begin{aligned} 6.16 \leq & \quad d & \leq 6.5 \\ 6.2 \times 10^8 \text{GeV} \geq & \quad \Lambda_{\rm IR} & \geq 1.0 \times 10^8 \text{GeV} \\ 1.6 \times 10^{-3} \leq & \quad \frac{m_{\rm direct}}{m_{\rm gauge}} & \leq 10^{-2} \end{aligned}$$

- Phenomenological Differences with conventional GMSB
  - Heavy Gravitino
  - Non-negligible FCNC
  - Presence of radion
  - . . .
- Work In Progress.

Conclusions

Final Word

- GMES is interesting as
  - An Alternative to GMSB
    - No need for traditional DSB
    - Averts gravitino constraints
    - Different phenomenology
  - Part of Susy w.o. Susy
    - No Susy flavor problem
    - Another class of realizations